

7184 North Park Drive Pennsauken, New Jersey 08109 (856) 486-1177 www.aerobiology.net

Hygenix, Inc
49 Woodside St.
Stamford , Connecticut 06902
Attn: Robert Brown

Project ID: WESTOVER SCHOOL

Date Collected: 10/27/2018
10/29/2018
10/29/2018
Project ID: 18041457

Project: STAMFORD P.S - WESTOVER SCHOOL

Condition of Sample(s) Upon Receipt: Acceptable

Project ID: 18041457

Page 1 of 5

1054 Spore Trap Analysis: SOP 3.8

Client Sample Number	26611659			26611653					
Sample Location		D225				D223			
Sample Volume (L)		75			75				
Lab Sample Number		18041457-	-001			18041457	-002		
Spore Identification	Raw Ct	spr/m³	% Ttl	In/Out	Raw Ct	spr/m³	% Ttl	In/Out	
basidiospores	-	-	-	-	1	53	20	-	
Cladosporium	2	107	67	-	-	-	-	-	
hyphal elements	-	-	-	-	1	53	20	-	
Pithomyces	-	-	-	-	1	53	20	_	
Smuts,Periconia,Myxomycetes	1	53	33	-	1	53	20	_	
Unknown	-	-	-	_	1	53	20	_	
		Debris Rating 2				Debris Rati	ng <b>2</b>		
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m³			Analyt	ical Sensitivi	ty: <b>13</b> s	pr/m³		
Comments									
Total *See Footnotes	3	160	~100%	-	5	267	~100%	-	

Client Sample Number	26611641				26611655				
Sample Location		D221				D219			
Sample Volume (L)		75				75			
Lab Sample Number		18041457-	003		18041457-004				
Spore Identification	Raw Ct	spr/m³	% Ttl	In/Out	Raw Ct	spr/m³	% Ttl	In/Out	
ascospores	-	-	-	-	1	53	12	-	
basidiospores	-	-	-	_	2	107	25	_	
Cladosporium	1	53	100	_	1	53	12	_	
Penicillium/Aspergillus group	-	-	-	-	4	213	50	-	
		Debris Ratir	ng <b>1</b>			Debris Rati	ng <b>1</b>		
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m³				Analyt	tical Sensitivit	y: <b>13</b> s	pr/m³	
Comments									
Total *See Footnotes	1	53	~100%	-	8	427	~100%	-	



7184 North Park Drive Pennsauken, New Jersey 08109 (856) 486-1177 www.aerobiology.net

Hygenix, Inc 49 Woodside St.

Stamford, Connecticut 06902

Attn: Robert Brown

Project: **STAMFORD P.S - WESTOVER SCHOOL** Condition of Sample(s) Upon Receipt: Acceptable

Date Collected: 10/27/2018
Date Received: 10/29/2018
Date Analyzed: 10/29/2018
Date Reported: 10/30/2018
Project ID: 18041457

Page 2 of 5

Client Sample Number	26611637					266116	346	
Sample Location		C213	C211					
Sample Volume (L)		75				75		
Lab Sample Number		18041457	-005			18041457	7-006	
Spore Identification	Raw Ct	spr/m³	% Ttl	In/Out	Raw Ct	spr/m³	% Ttl	In/Out
ascospores	2	107	9	-	4	213	13	-
basidiospores	1	53	4	-	2	107	6	-
Cladosporium	2	107	9	-	1	53	3	T -
hyphal elements	1	53	4	-	-	-	-	-
Penicillium/Aspergillus group	15	800	65	-	24	1280	75	-
Smuts,Periconia,Myxomycetes	1	53	4	-	1	53	3	-
Ulocladium	1	53	4	-	-	-	-	_
		Debris Rating 2				Debris Rat	ing <b>2</b>	
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m³			Analyt	ical Sensitivi	ity: <b>13</b> s	pr/m³	
Comments								
Total *See Footnotes	23	1227	~100%	-	32	1707	~100%	) -

Client Sample Number	26611635				2661163	6		
Sample Location	C209				C208			
Sample Volume (L)		75			75			
Lab Sample Number		18041457-	007			18041457-	800	
Spore Identification	Raw Ct	spr/m³	% Ttl	In/Out	Raw Ct	spr/m³	% Ttl	In/Out
ascospores	1	53	4	-	-	-	-	_
basidiospores	-	-	-	-	1	53	6	-
Chaetomium	-	-	_	-	1	53	6	-
hyphal elements	1	53	4	_	-	-	-	-
Penicillium/Aspergillus group	12	640	52	_	13	693	76	-
Rusts	1	53	4	_	-	-	-	-
Smuts,Periconia,Myxomycetes	7	373	30	-	1	53	6	-
Ulocladium	-	-	-	-	1	53	6	-
Unknown	1	53	4	_	-	-	-	-
	Debris Rating 3				Debris Rating 2			
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m³			Analytical Sensitivity: 13 spr/m³				
Comments								
Total *See Footnotes	23	1227	~100%	-	17	907	~100%	-



7184 North Park Drive Pennsauken, New Jersey 08109 (856) 486-1177 www.aerobiology.net

Hygenix, Inc 49 Woodside St.

Stamford, Connecticut 06902

Attn: Robert Brown

Project: **STAMFORD P.S - WESTOVER SCHOOL** Condition of Sample(s) Upon Receipt: Acceptable

Date Collected: 10/27/2018
Date Received: 10/29/2018
Date Analyzed: 10/29/2018
Date Reported: 10/30/2018
Project ID: 18041457

Page 3 of 5

Client Sample Number	26611664				26611645				
Sample Location		C207				C205			
Sample Volume (L)		75			75				
Lab Sample Number		18041457	-009			18041457	-010		
Spore Identification	Raw Ct	spr/m³	% Ttl	In/Out	Raw Ct	spr/m³	% Ttl	In/Out	
ascospores	1	53	5	-	1	53	9	-	
basidiospores	2	107	9	-	2	107	18	_	
Chaetomium	2	107	9	-	-	-	_	_	
Penicillium/Aspergillus group	8	427	36	-	7	373	64	-	
Smuts,Periconia,Myxomycetes	8	427	36	-	1	53	9	-	
Unknown	1	53	5	-	-	-	-	_	
		Debris Rating 2				Debris Rati	ng <b>2</b>		
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m³			<sup>3</sup> Analytical Sensitivity: <b>13</b> spr/m <sup>3</sup>			pr/m³		
Comments									
Total *See Footnotes	22	1173	~100%	-	11	587	~100%	-	

Client Sample Number	26611650				26611672				
Sample Location		C203				C201			
Sample Volume (L)		75				75			
Lab Sample Number		18041457-	011			18041457	'-012		
Spore Identification	Raw Ct	spr/m³	% Ttl	In/Out	Raw Ct	spr/m³	% Ttl	In/Out	
ascospores	1	53	11	-	-	-	-	-	
basidiospores	1	53	11	_	-	-	-	_	
Penicillium/Aspergillus group	3	160	33	_	4	213	67	-	
Smuts,Periconia,Myxomycetes	3	160	33	-	2	107	33	_	
Unknown	1	53	11	_	-	-	-	_	
		Debris Rating 2				Debris Rati	ing <b>1</b>		
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m³			<sup>13</sup> Analytical Sensitivity: <b>13</b> spr/m <sup>3</sup>			pr/m³		
Comments									
Total *See Footnotes	9	480	~100%	-	6	320	~100%	-	



Hygenix, Inc

49 Woodside St.

Stamford, Connecticut 06902

# **Certificate of Analysis** AIHA-LAP EMLAP# 102747

7184 North Park Drive Pennsauken, New Jersey 08109 (856) 486-1177 www.aerobiology.net

Date Collected: 10/27/2018 Date Received: 10/29/2018 Date Analyzed: 10/29/2018

Date Reported: 10/30/2018

Project ID: 18041457 Page 4 of 5

Attn: Robert Brown Project: STAMFORD P.S - WESTOVER SCHOOL Condition of Sample(s) Upon Receipt: Acceptable

Client Sample Number		2661169		2661167	'9			
Sample Location		TEACH W	STORAGE C					
Sample Volume (L)		75				75		
Lab Sample Number		18041457-	013			18041457-	014	
Spore Identification	Raw Ct	spr/m³	% Ttl	In/Out	Raw Ct	spr/m³	% Ttl	In/Out
ascospores	2	107	13	-	-	-	-	-
basidiospores	1	53	7	_	-	-	-	-
hyphal elements	-	-	-	-	1	53	7	-
Penicillium/Aspergillus group	11	587	73	-	12	640	80	-
Smuts,Periconia,Myxomycetes	1	53	7	_	-	-	-	-
Stachybotrys	-	-	-	_	2	107	13	-
		Debris Ratir			Debris Ratir	ng <b>1</b>		
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m³			Analytical Sensitivity: 13 spr/m³				
Comments								
Total *See Footnotes	15	800	~100%	_	15	800	~100%	-



7184 North Park Drive Pennsauken, New Jersey 08109 (856) 486-1177 www.aerobiology.net

Hygenix, Inc
49 Woodside St.
Stamford , Connecticut 06902
Attn: Robert Brown

Project ID: 10/27/2018
Date Collected: 10/27/2018
10/29/2018
Date Analyzed: 10/29/2018
Date Reported: 10/30/2018
Project ID: 18041457

Project: STAMFORD P.S - WESTOVER SCHOOL
Condition of Sample(s) Upon Receipt: Acceptable
Page 5 of 5

# **Footnotes and Additional Report Information**

### **Debris Rating Table**

1	Minimal (<5%) particulate present	Reported values are minimally affected by particulate load.
2	5% to 25% of the trace occluded with particulate	Negative bias is expected. The degree of bias increases directly with the percent of the trace that is occluded.
3	26% to 75% of the trace occluded with particulate	Negative bias is expected. The degree of bias increases directly with the percent of the trace that is occluded.
4	75% to 90% of the trace occluded with particulate	Negative bias is expected. The degree of bias increases directly with the percent of the trace that is occluded.
5	Greater than 90% of the trace occluded with particulate	Quantification not possible due to large negative bias. A new sample should be collected at a shorter time interval or other measures taken to reduce particulate load.

- 1. Penicillium/Aspergillus group spores are characterized by their small size, round to ovoid shape, being unicellular, and usually colorless to lightly pigmented. There are numerous genera of fungi whose spore morphology is similar to that of the Penicillium/Aspergillus type. Two common examples would be Paecilomyces and Acremonium. Although the majority of spores placed in this group are Penicillium, Aspergillus, or a combination of both. Keep in mind that these are not the only two possibilities.
- 2. Ascospores are sexually produced fungal spores formed within an ascus. An ascus is a sac-like structure designed to discharge the ascospores into the environment, e.g. Ascobolus.
- 3. Basidiospores are typically blown indoors from outdoors and rarely have an indoor source. However, in certain situations a high basidiospore count indoors may be indicative of a wood decay problem or wet soil.
- 4. The colorless group contains colorless spores which were unidentifiable to a specific genus. Examples of this group include Acremonium, Aphanocladium, Beauveria, Chrysosporium, Engyodontium microconidia, yeast, some arthrospores, as well as many others.
- 5. Hyphae are the vegetative mode of fungi. Hyphal elements are fragments of individual Hyphae. They can break apart and become airborne much like spores and are potentially allergenic. A mass of hyphal elements is termed the mycelium. Hyphae in high concentration may be indicative of colonization.
- 6. Dash (-) in this report, under raw count column means 'not detected (ND)'; otherwise 'not applicable' (NA).
- 7. The positive-hole correction factor is a statistical tool which calculates a probable count from the raw count, taking into consideration that multiple particles can impact on the same hole; for this reason the sum of the calculated counts may be less than the positive hole corrected total.
- 8. Due to rounding totals may not equal 100%.
- 9. Analytical Sensitivity for each spores is different for Non-viable sample when the spores are read at different percentage. Analytical Sensitivity is calculated as spr/m³ divided by raw count. spr/m³ = raw counts x (100/ % read) x (1000/Sample volume). If Analytical Sensitivity is 13 spr/m³ at 100% read, Analytical Sensitivity at 50% read would be 27 spr/m³, which is 2 times higher. Analytical Sensitivity provided on the report is based on an assumed 100% of the trace being analyzed.
- 10. Minimum Reporting Limits (MRL) for BULKS, DUSTS, SWABS, and WATER samples are a calculation based on the sample size and the dilution plate on which the organism was counted. Results are a compilation of counts taken from multiple dilutions and multiple medias. This means that every genus of fungi or bacteria recovered can be counted on the plate on which it is best represented.
- 11. If the final quantitative result is corrected for contamination based on the blank, the blank correction is stated in the sample comments section of the report.
- 12. The results in this report are related to this project and these samples only.
- 13. For samples with an air volume of < 100L, the number of significant figures in the result should be considered (2) two. For samples with air volumes between 100-999L, the number of significant figures in the result should considered (3) three. For example, a sample with a result of 55,443 spr/m³ from a 75L sample using significant figures should be considered 55,000. The same result of 55,443 from a 150L sample using significant figures should be considered 55,400 spr/m³.
- 14. If the In/Out ratio is greater than 100 times it is indicated >100/1, rather than showing the real value.

Terminology Used in Direct Exam Reporting

Conidiophores are a type of modified hyphae from which spores are born. When seen on a surface sample in moderate to numerous concentrations they may be indicative of fungal growth.

Suzanne S. Blevins, B.S., SM (ASCP) Laboratory Director

Suran 5. Bluing